

Chimera Experiment (2012)

- Omelchenko, Maistrenko, Hövel, Schöll:
PRL 106, 234102 (2011)
- Omelchenko, Riemenschneider, Hövel, Schöll:
PRE 85, 026212 (2012)

gekoppelte chaotische Abbildungen:

$$z_i^{t+1} = \underbrace{f(z_i^t)}_{(*)} + \frac{\sigma}{2P} \sum_{j=i-P}^{i+P} \underbrace{[f(z_j^t) - f(z_i^t)]}_{(**)} \quad i=1, \dots, N \quad (+) \text{ period. RB}$$


(*) lokale Dynamik: chaotisches Regime bei logistischer Abbildung

$$f(z) = az(1-z)$$

mit $a = 3,8$
(Lyapunov-Exponenten $\lambda = 0,931$)
(HA AG)

(**) nichtlokale Kopplung

- konstanter Kopplungskern
- Reichweite der nichtlokalen Kopplung P
und normierte Reichweite $r = \frac{P}{N}$
- Kopplungsstärke



Kontrollparameter - Ebene (σ, r)




FIG. 1 (color online). Regions of coherence for system (1) in the (r, σ) parameter plane with wave numbers $k = 1, 2$, and 3. Snapshots of typical coherent states z_i are shown in the insets. The color code inside the regions distinguishes different time periods of the states. The coherence-incoherence bifurcation (CIB) curve separates regions with coherent and incoherent dynamics. In the hatched and shaded (color) regions below CIB, two-cluster incoherent states exist. Completely synchronized chaotic states exist in the light hatched region bounded by the blowout bifurcation curve BB. Parameters: $a = 3.8$ and $N = 100$.

• $k=0$:
 zeitl. chaotisch,
 räumlich homogen
 (chaot. synchron)

• $k=1, 2, 3$:

Kohärenzzungen
 zeitlich periodisch : 2, 4, 8
 oberhalb CIB-Linie,
 ist das Profil "glatt"

CIB: coherence-incoherence
bifurcation

← Chaot. logistische Abb.





FIG. 2 (color online). Coherence-incoherence bifurcation for coupled chaotic logistic maps for fixed coupling radius $r = 0.32$ (black triangles in Fig. 1). For each value of the coupling parameter σ (decreasing from top to bottom, $\sigma = 0.43, 0.4, 0.32, 0.3, 0.2$, and 0.1 , respectively), snapshots (left columns) and space-time plots (right columns) are shown. Other parameters are as in Fig. 1.



Chaotische kontinuierliche
Dynamik: Rössler

FIG. 2. (Color online) Coherence regions in the (r, σ) parameter plane for $N = 100$ logistic maps (a) and Rössler systems (b), labeled by the wave number k . Gray scale (color code) inside the coherence regions distinguishes different periods and the coherence-incoherence bifurcation (CIB) curve separates regimes with coherent and incoherent dynamics. The light hatched region bounded by the blowout bifurcation curve BB refers to completely synchronized chaotic states. The insets in panel (a) display snapshots of typical coherent states. System parameters: panel (a) $a = 3.8$; panel (b) $a = 0.42, b = 2$, and $c = 4.0$. The vertical dashed lines refer to values of σ in Fig. 3. Triangles denote parameter values used to describe the CIB in Fig. 3.

Erster experimenteller Nachweis (2012)

Hagerstrom, Murphy, Roy, Hövel, Ouelchank, Schöll
 Nat. Phys. 8, 658 (2012)




Figure 1 | Experimental apparatus. **a**, Optical configuration. Polarization optics create a nonlinear relationship between the spatially dependent phase shift applied by the SLM and the intensity of the light falling on the camera. Feedback and coupling are implemented using a computer. **b,c**, Schematics of the 1D (**b**) and 2D (**c**) coupling configurations are shown. The site highlighted in white is updated based on the sites indicated in blue. As the elements are coupled diffusively to their neighbours within a range R in either one or two dimensions with periodic boundary conditions, the coupling is identical for all oscillators.




Figure 2 | Parameter space of the 1D and 2D CML systems. **a**, Parameter space of the 1D configuration ($N = 256$ elements). Blue corresponds to low entropy and periodicity; orange corresponds to high entropy and chaos. The dashed line indicates the critical coupling strength $\epsilon_c = 0.54$. There is a series of tongues containing profiles which are periodic in time and have spatial wavenumbers $K = 1, K = 2$ and $K = 3$ as indicated. **b**, Parameter space of the 2D configuration (128×128 lattice). There is a prominent $K = 1$ tongue, and two tongues with more complex spatial patterns that are not characterized by a wavenumber. **c**, Experimental and numerical realizations of the 1D system. In B, the incoherent region is highlighted in yellow. Labels A (coherence), B (chimera) and C (incoherence) show positions in the parameter space of **a**. **d**, Experimental and numerical realizations of the 2D system. In E, the incoherent region is highlighted.

Dynamik : $I(\phi) = \frac{1}{2} (1 - \cos \phi)$ "opt. Intensität"

Phase :
$$\phi_i^{t+1} = 2\pi a \left[I(\phi_i^t) + \frac{\epsilon}{2R} \sum_{j=-R}^R (I(\phi_{i+j}^t) - I(\phi_i^t)) \right]$$

$$= f(\phi_i^t) + \frac{\epsilon}{2R} \sum_{j=-R}^R (f(\phi_{i+j}^t) - f(\phi_i^t))$$

$\underbrace{\hspace{10em}}_{\pi a (1 - \cos \phi)}$


(1D-Ring)

ist chaotisch

kritische Kopplungsstärke ε_c

unterhalb der inkohärente Bereiche auftreten
(= Chimera-Zustände)

Profil:



- räumlich kontinuierlich : $N \rightarrow \infty$, $r = \frac{R}{N}$

$$\phi^{t+1}(x) = f(\phi^t(x)) + \frac{\varepsilon}{2r} \int_{x-r}^{x+r} dx' [f(\phi^t(x')) - f(\phi^t(x))] \quad \text{Rausgezogen}$$

- Betrachte $K=1$, Periode-2 Dynamik:

$$\phi^{t-j}(x) = (1-\varepsilon) f(\phi^j(x)) + \frac{\varepsilon}{2r} \int_{x-r}^{x+r} dx' f(\phi^j(x')) \quad ; j=0,1$$

- Bilde

$$\partial_x \phi^{t-j}(x) = (1-\varepsilon) \underbrace{f'(\phi^j(x))}_{\rightarrow \infty} \partial_x \phi^j + \frac{\varepsilon}{2r} \underbrace{[f(\phi^j(x+r)) - f(\phi^j(x-r))]}_{\text{vernachlässigen}}$$

bis Wendepunkt mit senkrechter Tangente

• Bilde $(\partial_x \phi^0, \partial_x \phi^1) = \left[(1-\varepsilon)^2 f'(\phi^0) f'(\phi^1) \right] \partial_x \phi^0 \cdot \partial_x \phi^1$

$(\Rightarrow) \quad 1 = (1-\varepsilon)^2 f'(\phi^0) f'(\phi^1) \quad | \quad f'(\phi) = \pi a \sin \phi$

$(\Rightarrow) \quad 0 = (1-\varepsilon)^2 (\pi a)^2 \sin \phi^0 \sin \phi^1 - 1 \stackrel{!}{=} G(x)$

• an den Diskontinuitätspunkten x^* gilt:

$\phi^0(x) = \phi^1(x) = \phi^* \hat{=} \text{keine 2-periodische Lösung, sondern Fixpunkt von } f$

$\rightarrow G(x) = (1-\varepsilon)^2 (\pi a)^2 \sin^2 \phi^*(x) - 1 = 0$

$$\varepsilon_c = 1 \quad (+) \quad \frac{1}{\pi a |\sin \phi^*|}$$

• " - " bestimmt den Threshold für ε

• $\phi^* \approx \phi_2$

$\Rightarrow \varepsilon_c \approx 0,55$




FIG. 5. Local map $f(\phi) = 2\pi a I(\phi) = \pi a(1 - \cos \phi)$ (red) and its twice iterate f^2 (blue) for $a = 0.85$.

The filled and open circles refer to the fixed points of f and the twice iterated f^2 .





FIG. 4. Simulations to illustrate scaling with system size of the incoherent regions for the 1D system. Snapshots for (a),(c),(e) $N = 256$, and (b),(d),(f) $N = 1024$. Decreasing coupling strength is used in panels (a),(b) $\epsilon = 0.44$, (c),(d) $\epsilon = 0.42$, (e),(f) $\epsilon = 0.40$. Other parameters: $a = 0.85$, $r = 0.41$. Transients of 1000 time steps were neglected. The insets show the enlarged left incoherence region for each snapshot.